

Wigner Molecules in Nanostructures

K. Jauregui, W. Häusler and B. Kramer

Physikalisch–Technische Bundesanstalt Braunschweig

Bundesallee 100, 3300 Braunschweig F. R. G.

February 1, 2008

Abstract

The one- and two- particle densities of up to four interacting electrons with spin, confined within a quasi one-dimensional “quantum dot” are calculated by numerical diagonalization. The transition from a dense homogeneous charge distribution to a dilute localized Wigner-type electron arrangement is investigated. The influence of the long range part of the Coulomb interaction is studied. When the interaction is exponentially cut off the “crystallized” Wigner molecule is destroyed in favor of an inhomogeneous charge distribution similar to a charge density wave .

PACS. 71.45.Lr – Charge-density-wave systems.

PACS. 73.20.Dx – Electron states in low-dimensional structures.

PACS. 73.20.Mf – Collective excitations.

Single charges dominate the electronic properties of submicron structures at low temperatures. Due to small capacitances the charging energy associated with the addition of one electron into a given structure can exceed the thermal energy. Two recently discovered important phenomena can be explained by single electron charging effects: the Coulomb blockade of the dc-current through small tunnel junctions [1, 2], and the periodic oscillations of the conductance of quantum dots [3, 4].

In contrast to metallic systems semiconductor nano-structures allow to reduce the number of electrons in a quantum dot by changing a gate voltage. Optical absorption experiments were performed on quantum dots which contain only $N = 2 \dots 4$ electrons [5]. For very low electron densities additional effects in the transport properties can be expected due to the increasing importance of the Coulomb interaction [6] since the electrons tend to “crystallize” into an inhomogeneous ground state [7]. Besides Coulomb effects, linear [8] and non-linear [9] transport experiments show fine structure in the current voltage characteristics that can be traced back to the granularity of the charge density distribution and to the lowest collective excitations [10], respectively.

In this paper, we demonstrate that for a quasi-one dimensional (1D) system containing a few electrons the charge density $\varrho(x)$ becomes inhomogeneous for sufficiently large mean electron distance r_s . Our results indicate that there are two consecutive regimes of charge localization. First, $\varrho(x)$ starts to become inhomogeneous with increasing r_s . The corresponding energetically lowest excitations are associated with changes in symmetry and total spin [11]. When r_s is further increased, $\varrho(x)$ vanishes almost completely between well separated maxima. In analogy with the Wigner lattice, we denote this limiting configuration in a finite system as a “Wigner molecule”. Two classes of elementary excitations can be identified in this limit: tunneling between different permutational arrangements of the electrons and vibrations of the charge density [6, 11].

We consider N interacting electrons including spin, confined in a 1D square well potential of finite length L . The height of the potential was assumed to be finite but large. The Hartree-Fock approximation is known [12] to overestimate the ferromagnetic state and cannot reproduce the true ground state which is antiferromagnetic in 1D [13]. We calculate eigenvalues and eigenstates of the Hamiltonian exactly.

The kinetic energy

$$E_H \left(\frac{a_B}{L} \right)^2 \sum_{n,\sigma} \epsilon_n c_{n,\sigma}^\dagger c_{n,\sigma} \equiv E_H \left(\frac{a_B}{L} \right)^2 H_0 \quad (1)$$

and the Coulomb interaction

$$E_H \left(\frac{a_B}{L} \right) \sum_{\substack{n_1 \dots n_4 \\ \sigma, \sigma'}} V_{n_4 n_3 n_2 n_1} c_{n_4 \sigma}^\dagger c_{n_3 \sigma'}^\dagger c_{n_2 \sigma'} c_{n_1 \sigma} \equiv E_H \left(\frac{a_B}{L} \right) H_I \quad (2)$$

scale like $1/L^2$ and $1/L$, respectively. Thus the latter dominates at large L . The natural energy and length scales are given by the Hartree $E_H = e^2/a_B$, and the Bohr radius, $a_B = \varepsilon \hbar^2 / m e^2$ (ε relative dielectric constant, m effective electron mass). $c_{n,\sigma}^\dagger$ creates a particle with energy ϵ_n ($= (n\pi)^2/2$ for an infinitely high well) and spin σ . $V_{n_4 n_3 n_2 n_1}$ is the matrix element of the interaction potential $V(x, x') = e^2 / \varepsilon \sqrt{(x - x')^2 + \lambda^2}$. The cutoff at short distances simulates a small transversal spread $\lambda \ll L$ of the wave functions and leaves the $V_{n_4 n_3 n_2 n_1}$ finite. In the calculations we assumed $\lambda/L = 2 \cdot 10^{-4}$. The eigenvalues of $H = E_H(a_B/L)[(a_B/L)H_0 + H_I]$ depend only weakly on λ when $\lambda \ll 1$ [6]. To investigate the effect of the long range part of the interaction, in some of the calculations an exponential cutoff at large distances was introduced. No charge neutralizing background was taken into account since it is not expected to modify the results for fixed N qualitatively.

The Hamiltonian matrix in the basis of the Slater determinants was diagonalized numerically. The latter were constructed from the single electron states $\varphi_n(x)\chi_\sigma$ where φ_n is a spatial function and χ_σ a spinor with $\sigma = \downarrow, \uparrow$. Considering $1 \leq n \leq M$ leads to a matrix of the rank $R = \binom{2M}{N}$ ($R < 1.5 \cdot 10^4$, for $M = 10 \dots 17$). We used Lanczos procedures when $R > 3 \cdot 10^3$.

The one- and two-particle densities were determined from the normalized eigenfunctions $\psi(x_1, \dots, x_N) = \sum_{\nu=1}^R (b_\nu / \sqrt{N!}) \det\{\varphi_{n_i}(x_j)\chi_{\sigma_i}\}$, where ν denotes the configurations $\{n_1\sigma_1, \dots, n_N\sigma_N\}$. The coefficients b_ν can be chosen real. In second quantization the one-particle density is given by

$$\varrho(x) = \sum_{\sigma} \langle \psi_0^{(N)} | \Psi_{\sigma}^\dagger(x) \Psi_{\sigma}(x) | \psi_0^{(N)} \rangle, \quad (3)$$

where $|\psi_0^{(N)}\rangle = \sum_{\nu=1}^R b_\nu^0 |\nu\rangle$ is the N -electron ground state. $\Psi_{\sigma}^\dagger(x) = \sum_{n=1}^M \varphi_n(x) c_{n,\sigma}^\dagger$ creates an electron at position x with spin σ and $|\nu\rangle = c_{n_1\sigma_1}^\dagger \dots c_{n_N\sigma_N}^\dagger |0\rangle$ is a noninteracting Slater determinant. Straightforwardly we have

$$\varrho(x) = \sum_{\substack{\nu=1 \\ \nu'=1}}^R b_{\nu}^0 b_{\nu'}^0 \sum_{\substack{n, n' \\ \sigma}} \varphi_n(x) \varphi_{n'}(x) \langle \nu' | c_{n', \sigma}^{\dagger} c_{n, \sigma} | \nu \rangle \quad . \quad (4)$$

The development of maxima and minima in $\varrho(x)$ with increasing mean electron distance $r_s \equiv L/(N-1)$ is shown in figure 1 for $N = 3, 4$. The finite height of the square well potential was assumed to be proportional to $1/L^2$ such that the number of non-interacting bound states was independent of L . This implies a non-vanishing but small $\varrho(x)$ outside the box.

We distinguish three characteristic regimes of electron densities. For small distances, $r_s \lesssim 0.1a_B$, the Coulomb interaction perturbs $E_H(a_B/L)^2 H_0$ only weakly such that $\varrho(x)$ is dominated by the lowest occupied single particle states. This explains the minimum at $x = 0$ (Fig.1). $\varrho(x)$ changes qualitatively at $r_s \approx a_B$. A structure consisting of N peaks emerges. The “critical” length for this transition is of the same order as found in [11] for the transformation from an almost non-interacting energy spectrum to the multiplet spectrum dominated by interaction. When r_s increases further, say $r_s \gtrsim 100a_B$, $\varrho(x)$ vanishes almost completely in finite regions between the maxima indicating a fully established Wigner molecule. In this limit the ground state energy can be approximated by that of a chain of elementary charges at equal distances r_s .

In order to investigate the influence of the long range part of the Coulomb interaction, we introduced an exponential cutoff $V(x, x') \propto e^{-\alpha|x-x'|}/\sqrt{(x-x')^2 + \lambda^2}$, where $1/\alpha$ is the cutoff distance. The inset of figure 1 shows $\varrho(x)$ for $N = 3$ and $\alpha = 10a_B^{-1}$. Despite the extremely short range of the interaction pronounced maxima are observed. However, in contrast to $\alpha = 0$ the density between the maxima is much higher even for very large r_s . The inset of figure 1 rather resembles a charge density wave [14]. We conclude that the long range part of the Coulomb interaction is essential for the charge condensation and for long range density-density correlations. The ground state energy for $\alpha > r_s^{-1}$ does not converge towards the Coulomb energy of N elementary point charges for $L \rightarrow \infty$.

For large r_s ($r_s \gg a_B$) the spectrum consists of multiplets [15]. Their energetic separation Ω correspond to vibrational excitations. Their internal splitting Δ is due to tunnelling through the Coulomb barrier between adjacent permutational arrangements of the electrons [11]. Asymptotically, Ω decreases as $r_s^{-\gamma}$. Power law behavior is also obtained for $\alpha > 0$. However, the exponent γ changes from $\gamma \lesssim 1.5$ ($\alpha = 0$) to $\gamma \approx 2$ ($\alpha > 0$) (figure 2). The vibrational excitations are independent of the symmetry of the spatial part of the N -

electron wave function and of the total spin. In analogy to the argument for $\alpha = 0$ leading to $1 \leq \gamma \leq 1.5$ [11], the almost flat two-particle potential between adjacent charge density peaks for $\alpha \gg r_s^{-1}$ leads to $\gamma = 2$. Each electron is practically moving freely within a box of length r_s . On the other hand, Δ decreases roughly exponentially with increasing L , even for $\alpha > 0$ (inset of figure 2). However, Δ is larger and the decrease is less rapid when r_s is increased than for $\alpha = 0$. The enhancement results from the reduced thickness of the screened Coulomb barrier such that the corresponding tunnelling integral ($\propto \Delta$) increases. The modified dependence on r_s is due to the fact that only for $r_s < \alpha^{-1}$ the thickness of the interaction barrier varies with r_s . We find that the tunnelling excitations can be described in terms of correlated, localized basis functions even for $\alpha > 0$. However, in order to obtain similar quantitative agreement with the numerically determined eigenvalues as for $\alpha = 0$ a larger mean electron distance r_s is required to fulfill $\Delta \ll \Omega$. As a consequence the characteristic length for the transition from almost non-interacting to the interaction dominated behavior increases with increasing α .

The pronounced correlations between the electrons in our model are also observed in the pair correlation function of the N -electron state $|\psi^{(N)}\rangle$,

$$\varrho_c(x, x') = \sum_{\sigma, \sigma'} \langle \psi^{(N)} | \Psi_{\sigma}^{\dagger}(x) \Psi_{\sigma'}^{\dagger}(x') \Psi_{\sigma'}(x') \Psi_{\sigma}(x) | \psi^{(N)} \rangle . \quad (5)$$

ϱ_c is related to the density-density correlation function ϱ_2 via $\varrho_2(x, x') = \varrho(x)\delta(x - x') + \varrho_c(x, x')$. Physically, ϱ_c is the probability to find an electron at position x when another electron sits at x' ($\int dx \int dx' \varrho_c(x, x') = N(N-1)$). Figure 3 shows a contour plot of ϱ_c for the $N = 3$ ground state ($S = 1/2$) with $L = 9.5a_B$. Practically the same result is obtained for the spin polarized excited state ($S = 3/2$). The difference between ϱ_c of the ground state and the spin polarized state is less than 1%, independent of x , and x' .

It is known that the Hartree Fock approximation overestimates the tendency of electrons with parallel spins to avoid each other and underestimates the correlations between electrons with antiparallel spins. Therefore large differences between correlation functions for different spin configurations are expected in this approximation. Recently Pfannkuche et al. demonstrated this property explicitly for three electrons in a two dimensional parabolic potential [16]. The weak sensitivity of the pair correlation function on S shows that highly correlated electrons in a “slim” quantum dot cannot be treated within molecular field approximation.

In summary, we have analyzed one- and two-particle densities for the lowest eigenstates

of $N = 2, 3, 4$ electrons confined in a quasi-1D square well. Three different regimes were identified. At high electron densities the Coulomb interaction is only a weak perturbation. Between $a_B^{-1} \gtrsim r_s^{-1} \gtrsim 10^{-2} a_B^{-1}$ the electrons tend to localize. Simultaneously the eigenvalue spectrum develops a multiplet structure that is dominated by the interaction. Both, the inhomogeneity of the charge density and the multiplet character of the spectrum are not destroyed when the long range part of the interaction is removed. However, there are quantitative changes in the asymptotic behavior of tunneling and vibrational excitations at low densities. For the lowest densities (equivalent to $r_s \gtrsim 100 a_B$) the formation of a *Wigner molecule* is observed. In this limit where the long range part of the interaction is crucial, $\rho(x)$ vanishes within finite intervals between the maxima. The structure in the density-density correlation function becomes more pronounced with increasing influence of the interaction but is less sensitive to total electron spin than expected within Hartree Fock approximation. It is worth remarking that the best numerical estimate to our knowledge for the critical value of r_s for the formation of a Wigner crystal in 3D is also $r_s^{\text{cr}} \approx 100 a_B$ [17].

Experiments are frequently performed on 2D quantum dots that are based on GaAs-AlGaAs heterostructures. In many of these the electron density corresponds to the intermediate regime. For the geometry and the electron numbers in typical quantum dots [3, 4] $r_s \approx 3 a_B$ (area of the dot $\approx 10^5 \text{ nm}^2$, number of electrons $\approx 10^2$, effective mass $\approx 0.07 m_e$, dielectric constant ≈ 10). Even for this relatively high electron density in comparison with more recent experiments [5] the charge density distribution cannot be expected to be homogeneous if we assume that the qualitative features of the electronic properties discussed above apply also to the 2D case. In small quantum dots that are weakly coupled to leads and that contain only a finite number of electrons, like the experimentally investigated double barrier systems [3], a granular charge density should lead to characteristic features in linear transport properties as a function of the gate voltage V_G . A variation of V_G not only changes the charge density or the electron number inside the dot but probably simultaneously its shape. In [4] the gate electrode is positioned besides the dot region. For this geometry the influence of V_G on the shape of the dot area is obvious. The resonance condition for the appearance of a conductance peak in linear transport $E_F = E_{N+1} - E_N$ may be fulfilled more than once for given N when V_G is increased. Here E_F is the chemical potential of the leads (where no electron-electron interaction is assumed) and E_N the N -electron *ground state* energy of the quasi-isolated dot. Such a model could account for the experimentally

observed fine structure in the conductance peaks in the *linear* transport regime reported in [8] which seem to be presently not yet very well understood.

This work was partly supported by grants of the Deutsche Forschungsgemeinschaft (AP 47/1-1) and EEC (Science Contract No.SCC*– CT90–0020).

Figure 1: (a) One-particle density $\varrho(x)$ for $N = 3$ and (b) $N = 4$ for different L ($0.1a_B \leq L \leq 945a_B$, $M = 13$). The normalisation is such that $\int dx \varrho(x) = N$. When $L \gtrsim 1a_B$ N peaks begin to emerge. For $L \gtrsim 100a_B$ the peaks are well separated. Inset: $\varrho(x)$ for a pair potential with an exponential cutoff ($\alpha = 10a_B^{-1}$) for $N = 3$ and $L = 0.1a_B$ (dotted line); $L = 9.5a_B$ (dashed-dotted line); $L = 472a_B$ (dashed line); $L = 1417a_B$ (solid line) ($M = 11$). The minima at large system lengths are less pronounced as for $\alpha = 0$.

Figure 2: Influence of a pair potential on the energy difference between the lowest two multiplets Ω (vibrational excitation). The ratio $\Omega(\alpha = 10a_B^{-1})/\Omega(\alpha = 0)$ is plotted versus the particle distance r_s for $N = 2$ ($\times$) and $N = 3$ ($\circ$). ($M = 11$). Inset: logarithm of the energy difference Δ between the ground state and the first excited state within the lowest multiplet versus system length L for $N = 2$, $\alpha = 0$ ($+$) and $\alpha = 10a_B^{-1}$ ($\circ$); $N = 3$, $\alpha = 0$ ($\times$) and $\alpha = 10a_B^{-1}$ (Y). ($M = 11$).

Figure 3: Contour plot of the pair correlation function $\varrho_c(x, x')$ (cf eq. 5) for the $N = 3$ ground state ($S = 1/2$) at $L = 9.5a_B$ ($M = 13$).

References

- [1] D. V. Averin, K. K. Likharev in : *Quantum Effects in Small Disordered Systems*, ed. by B. L. Altshuler, P. A. Lee, R. A. Webb (Elsevier, Amsterdam, 1991);
G. Schön, A. D. Zaikin, Phys. Rep. **198**, 237 (1990);
H. Grabert, G. L. Ingold, M. H. Devoret, D. Esteve, H. Pothier, C. Urbina, Z. Phys. B**84**, 143 (1991);
T. A. Fulton and G. J. Dolan, Phys. Rev. Lett. **59**, 109 (1987).
- [2] Special Issue on *Single Charge Tunneling* edited by H. Grabert, Z. Phys. B**85**, 317–468 (1991).
- [3] U. Meirav, M. A. Kastner, S. J. Wind, Phys. Rev. Lett. **65**, 771 (1990);
L. P. Kouwenhoven, N. C. van der Vaart, A. T. Johnson, W. Kool, C. J. P. M. Harmans, J. G. Williamson, A. A. M. Staring, C. T. Foxon in Ref. [2], p. 367.
- [4] M. A. Kastner, Rev. Mod. Phys. **64**, 849 (1992).
- [5] Ch. Sikorski, U. Merkt, Phys.Rev.Lett. **62**, 2164 (1989);
B. Meurer, D. Heitmann, K. Ploog, Phys. Rev. Lett. **68**, 1371 (1992).
- [6] W. Häusler, B. Kramer, J. Mašek in Ref. [2], p. 435.
- [7] Wigner E, Phys. Rev. **46**, 1002 (1934).
- [8] L. P. Kouwenhoven: PhD thesis, University of Delft (1992), which contains a collection of papers on this subject.
- [9] A. T. Johnson, L. P. Kouwenhoven, W. de Jong, N. C. van der Vaart, C. J. P. M. Harmans, and C. T. Foxon, Phys. Rev. Lett. **69**, 1592 (1992)
- [10] W. Pfaff, D. Weinmann, W. Häusler, B. Kramer, U. Weiss, to be published.
- [11] W. Häusler, B. Kramer, to be published in Phys. Rev. B.
- [12] D. C. Mattis, Physics **1**, 183 (1964).
- [13] E. Lieb, D. Mattis, Phys. Rev. **125**, 164 (1962).
- [14] I. M. Tsidil'kovskii, Sov. Phys. Usp. **30**, 676 (1987).

- [15] There are 2^N (equal to the dimension of the spin Hilbert space) states per multiplet, differing in total spin and in symmetry with respect to permutations of the spatial space coordinates of the N -electron wave function [11].
- [16] D. Pfannkuche, V. Gudmundsson, P. A. Maksym, Phys. Rev. B **47**, 2244 (1993).
- [17] D. M. Ceperley, B. J. Adler, Phys. Rev. Lett. **45**, 566 (1980)